

EAU GUIDELINES ON UROLOGICAL TRAUMA

(Limited text update March 2018)

N.D. Kitrey (Chair), N. Djakovic, F.E. Kuehhas, N. Lumen,
E. Serafetinidis, D.M. Sharma
Guidelines Associates: Y. Abu-Ghanem, A. Sujenthiran

Introduction

Genito-urinary trauma is seen in both sexes and in all age groups, but is more common in males. Traumatic injuries are classified according to the basic mechanism into **penetrating** and **blunt**.

Penetrating trauma is further classified according to the velocity of the projectile:

1. High-velocity projectiles (e.g. rifle bullets - 800-1000 m/sec).
2. Medium-velocity (e.g. handgun bullets - 200-300 m/sec).
3. Low-velocity items (e.g. knife stab).

High-velocity weapons inflict greater damage as the bullets transmit large amounts of energy to the tissues, resulting in damage to a much larger area than the projectile tract itself. In lower velocity injuries, the damage is usually confined to the tract of the projectile.

Blast injury is a complex cause of trauma, it commonly includes both blunt and penetrating trauma, and may also be accompanied by burn injuries.

Initial evaluation and management

The first priority is stabilisation of the patient and treatment of associated life-threatening injuries. A direct history is obtained from the patient (if conscious) or from witnesses/emergency personnel (if patient is unconscious and/or seriously injured).

In penetrating injuries, assess size of the weapon in stabbings, and the type and calibre of the weapon used in gunshot wounds. The medical history should be as detailed as possible. It is important to recognise the high risk of hepatitis B and C infection in trauma patients and take appropriate precautions. In any penetrating trauma, tetanus vaccination should be considered according to the patient's vaccination history and nature of the wound.

Renal Trauma

Renal injuries are associated with young age and male gender, the incidence is approximately 4.9 per 100,000 of the population. The most commonly used classification system is that of the American Association for the Surgery of Trauma. This validated system has clinical relevance and helps to predict the need for intervention. It also predicts morbidity after blunt or penetrating injury and mortality after blunt injury.

Diagnostic evaluation

Recommendations	Strength rating
Assess haemodynamic stability upon admission.	Strong
Record past renal surgery, and known pre-existing renal abnormalities (ureteropelvic junction obstruction, large cysts, lithiasis).	Strong
Test for haematuria in a patient with suspected renal injury.	Strong

<p>Perform a contrast enhanced CT scan in blunt trauma patients with;</p> <ul style="list-style-type: none"> • visible haematuria; • non-visible haematuria with haemodynamic instability; • a history of rapid deceleration injury and/or significant associated injuries; • penetrating abdominal or lower thoracic injury. 	Strong
<p>Perform CT delayed phase images in case of haemodynamic stability.</p>	Strong

Management

Recommendations	Strength rating
<p>Manage stable patients with blunt renal trauma conservatively with close monitoring of vital signs.</p>	Strong
<p>Manage isolated grade 1-3 stab and low-velocity gunshot wounds in stable patients, expectantly.</p>	Strong
<p>Use angioembolisation for active renal bleeding if there are no other indications for immediate laparotomy.</p>	Strong
<p>Proceed with renal exploration in the presence of:</p> <ul style="list-style-type: none"> • persistent haemodynamic instability; • expanding or pulsatile peri-renal haematoma; • grade 5 vascular injury; • exploration for associated injuries. 	Strong
<p>Attempt renal reconstruction if haemorrhage is controlled and there is sufficient viable renal parenchyma.</p>	Weak

Figure 1: Evaluation of blunt renal trauma in adults





Figure 2: Evaluation of penetrating renal trauma in adults



* Suspected renal trauma results from reported mechanism of injury and physical examination.

† Renal imaging: CT scans are the gold standard for evaluating blunt and penetrating renal injuries in stable patients. In settings where CT is not available, the urologist should rely on other imaging modalities (IVP, angiography, radiographic scintigraphy, MRI).

‡ Renal exploration: Although renal salvage is a primary goal for the urologist, decisions concerning the viability of the organ and the type of reconstruction are made during the operation.

CT = computed tomography; Ht = haematocrit;
IVP = intravenous pyelography.

Post-operative care, follow-up and complications

Recommendations	Strength rating
Repeat imaging in case of fever, worsening flank pain, or falling haematocrit.	Strong
Follow-up approximately three months after major renal injury with: <ul style="list-style-type: none">• physical examination;• urinalysis;• individualised radiological investigation including nuclear scintigraphy;• serial blood pressure measurements;• renal function tests.	Weak

Iatrogenic renal injuries

- Iatrogenic renal injuries are procedure-dependent (1.8-15%).
- Significant injury requiring intervention is rare; the most common injuries are vascular.
- Renal allografts are more susceptible.
- Injuries occurring during surgery should be rectified immediately.
- Symptoms suggestive of a significant injury require investigation.

The recommendations for iatrogenic renal injuries are the same as those for follow-up.

Ureteral Trauma

Ureteral injuries are quite rare - most are iatrogenic. They are often missed intra-operatively, usually involve the lower ureter, and may result in severe sequelae. Risk factors include advanced malignancy, prior surgery or irradiation - i.e. conditions which alter the normal anatomy. Pre-operative prophylactic stents do not prevent ureteral injury, but may

assist in its detection. External ureteral trauma usually accompanies severe abdominal and pelvic injuries. Gunshot wounds account for the majority of penetrating ureteral trauma, while motor vehicle accidents account for most blunt injuries.

Diagnostic evaluation

- A high index of suspicion of ureteral injury should be maintained as the majority of cases are diagnosed late, predisposing the patient to pain, infection, and renal function impairment.
- Haematuria is an unreliable indicator.
- Extravasation of contrast material in computed tomography (CT) is the hallmark sign of ureteral trauma.
- In unclear cases, a retrograde or antegrade urography is required for confirmation.

Management

Recommendations	Strength rating
Visually identify the ureters to prevent ureteral trauma during abdominal and pelvic surgery.	Strong
Beware of concomitant ureteral injury in all abdominal penetrating trauma, and in deceleration-type blunt trauma.	Strong
Use pre-operative prophylactic stents in high-risk cases.	Strong

The type of repair procedure depends on the site of the injury (Table 2), and should follow the principles outlined in Table 3.

Site of injury	Reconstruction options
Upper ureter	Uretero-ureterostomy Transuretero-ureterostomy Uretero-calycostomy
Mid ureter	Uretero-ureterostomy Transuretero-ureterostomy Ureteral re-implantation and a Boari flap
Lower ureter	Ureteral re-implantation Ureteral re-implantation with a psoas hitch
Complete	Ileal interposition graft Autotransplantation

Debridement of necrotic tissue
Spatulation of ureteral ends
Watertight mucosa-to-mucosa anastomosis with absorbable sutures
Internal stenting
External drain
Isolation of injury with peritoneum or omentum

Bladder Trauma

Bladder injuries can be due to external (blunt or penetrating) or iatrogenic trauma. Iatrogenic trauma is caused by external laceration or internal perforation (mainly during transurethral resection of the bladder). Blunt bladder injuries are strongly associated with pelvic fractures. Bladder injuries are classified as extraperitoneal, intraperitoneal or combined.

Diagnostic evaluation

Clinical signs and symptoms

External trauma

- Cardinal sign: visible haematuria.
- Others: abdominal tenderness, inability to void, bruises over the suprapubic region, and abdominal distension (in case of urinary ascites).
- Penetrating bladder injury: entrance and exit wounds in lower abdomen or perineum.
- Bloody urethrorrhagia: suspect concomitant urethral injury.

Iatrogenic trauma

- External perforation: extravasation of urine, visible laceration, clear fluid in the surgical field, appearance of the bladder catheter, and blood and/or gas (in case of laparoscopy) in the urine bag.
- Internal perforation: fatty tissue or bowel between detrusor muscle fibres, inability to distend the bladder, low return of irrigation fluid and/or abdominal distension.
- Post-operative symptoms of unrecognised bladder perforation: haematuria, lower abdominal pain, abdominal distension, ileus, peritonitis, sepsis, urine leakage from the wound, decreased urinary output, and increased serum creatinine.

Imaging – Cystography and Cystoscopy

Recommendations	Strength rating
Perform cystography in case of suspected iatrogenic bladder injury in the post-operative setting.	Strong
Perform cystography in the presence of visible haematuria and pelvic fracture.	Strong

Perform cystography with active retrograde filling of the bladder with dilute contrast.	Strong
Use cystoscopy to rule out bladder injury after suburethral sling procedure by the retropubic route.	Strong
Do not perform cystography to assess bladder wall healing after repair of a simple injury in a healthy patient.	Weak
Perform cystography to assess bladder wall healing after repair of a complex injury or in case of risk factors for wound healing.	Strong

Management

- Surgical repair (two-layer vesicorrhaphy).
- Conservative management (urinary catheter).

Recommendations	Strength rating
Manage a blunt extraperitoneal bladder injury operatively in cases of bladder neck involvement and/or associated injuries that require surgical intervention.	Strong
Manage uncomplicated blunt extraperitoneal bladder injury conservatively.	Weak
Manage intraperitoneal injuries caused by blunt trauma by surgical exploration and repair.	Strong
Manage small uncomplicated iatrogenic intraperitoneal bladder injuries conservatively.	Weak

Urethral Trauma

- Injuries to the anterior urethra (AU) are caused by trauma during sexual intercourse (associated with penile fracture), penetrating trauma, placement of penile constriction bands, and from iatrogenic trauma e.g. endoscopic instruments, catheterisation.
- Injuries to the posterior urethra (PU) occur with pelvic fractures, mostly as a result of motor vehicle accidents.
- The male PU is injured in 4-19% of pelvic fractures, and the female urethra in 0-6% of all pelvic fractures.
- The combination of straddle fractures with diastasis of the sacroiliac joint has the highest risk of urethral injury.
- Injuries can vary from simple stretching to partial rupture to complete disruptions.
- Urethral injuries in women are rare.

Diagnostic evaluation

- Blood at the external urethral meatus is the most common clinical sign, and indicates the need for further diagnostic work up.
- Although non-specific, haematuria on a first voided specimen may indicate urethral injury. The amount of urethral bleeding correlates poorly with the severity of injury.
- Pain on urination or inability to void may indicate disruption.
- Blood at the vaginal introitus is present in more than 80% of female patients with pelvic fractures and co-existing urethral injuries.
- Rectal examination may reveal a "high riding" prostate. However, this is an unreliable finding. Blood on the examination finger is suggestive of a rectal injury associated with pelvic fracture.
- Urethral bleeding or urinary extravasation can cause penile and scrotal swelling and haematoma.

- Retrograde urethrography is the gold standard for evaluating urethral injury and urethral catheterisation should be avoided until the urethra is imaged.
- In an unstable patient, however, an attempt can be made to pass a urethral catheter (gently, by someone with urological experience). If this is not possible, a suprapubic catheter is inserted and a retrograde urethrogram is performed later.
- In females, urethroscopy may be an important adjunct for the identification and staging of urethral injuries.

Management

Urethral trauma

Recommendations	Strength rating
Evaluate urethral injuries with flexible cystoscopy and/or retrograde urethrography.	Strong
Treat blunt anterior urethral injuries by suprapubic diversion.	Strong
Treat partial posterior urethral ruptures by urethral or suprapubic catheterisation.	Strong
Perform early endoscopic re-alignment when feasible.	Weak
Manage complete posterior urethral disruption with suprapubic diversion and deferred (at least three months) urethroplasty.	Strong

Iatrogenic urethral trauma

Recommendations	Strength rating
Provide appropriate training to reduce the risk of traumatic catheterisation.	Strong
Keep duration of catheterisation to a minimum.	Strong

Genital Trauma

Of all genito-urinary injuries, one-third to two-thirds involve the external genitalia and are much more common in males due to anatomical differences, increased frequency of road traffic accidents, physical sports, violent crime, and war-fighting. Eighty percent is blunt trauma, 20% is due to penetrating injuries.

Diagnostic evaluation

- Urinalysis should be performed.
- Visible- and/or non-visible haematuria requires a retrograde urethrogram in males, whilst cystoscopy should be considered in females.
- In women with genital injuries and blood at the vaginal introitus, further gynaecologic investigation to exclude vaginal injury is required.
- In cases of suspected sexual abuse gynaecologic and forensic support and advice is necessary. The emotional situation and privacy of the patient must be respected.

Blunt penile trauma

Usually results from trauma to the erect penis during sexual intercourse or masturbation.

Penile fracture

- Sudden cracking or popping sound, pain and immediate detumescence.
- Local swelling of the penile shaft is seen and this may extend to the lower abdominal wall.
- The rupture of the tunica may be palpable.
- Thorough history and examination confirms diagnosis.
- Imaging ultrasound (US) or magnetic resonance imaging (MRI) may be useful.

Management

- Subcutaneous haematoma, without associated rupture of the cavernosal tunica albuginea does not require surgical intervention. Non-steroidal analgesics and ice-packs are recommended.
- In penile fracture, early surgical intervention with closure of the tunica albuginea is recommended.
- Intra-operative flexible cystoscopy is useful to diagnose urethral injury and to further localise tunical damage.
- Conservative management of penile fracture is not recommended.

Penetrating penile trauma

- Rarely seen in isolation.
- Due to gunshot/knife injury, animal or human bites, assault and industrial or self-inflicted mutilation.
- Non-operative management is recommended in small superficial injuries with intact Buck's fascia.
- More significant injuries require surgical exploration and debridement of necrotic tissue.
- In extended injuries of the penis, primary alignment of the disrupted tissues may allow for acceptable healing because of the robust penile blood supply.
- In avulsion of the penis, resuscitate the patient and attempt re-implantation of the penis (if not too badly damaged) - ideally microsurgically.

Blunt scrotal trauma

- May result in testicular dislocation, haematocoele, testicular rupture and/or scrotal haematoma.
- Dislocation of the testicle is rare. Treat by manual replacement and secondary orchidopexy. If manual repositioning cannot be performed, immediate orchidopexy is indicated.
- If haematocoele is smaller than three times the size of the contralateral testis – conservative management.

- If large haematocele - explore.
- If testicular rupture suspected, explore, evacuate clot and any necrotic testicular tubules and close the tunica albuginea.

Penetrating scrotal trauma

- Surgical exploration with conservative debridement of nonviable tissue.
- Primary reconstruction of testis and scrotum can be performed in most cases.
- In complete disruption of the spermatic cord, re-alignment without vaso-vasostomy may be considered.
- In extensive destruction of the tunica albuginea, mobilisation of a free tunica vaginalis flap can be performed for testicular closure.
- If reconstruction cannot be achieved, orchiectomy is indicated.
- In improvised explosive device blast injury, the extensive loss of genital tissue often requires complex and staged reconstructive surgical procedures.

Genital trauma in females

- In blunt trauma to the external genitalia, imaging studies of the pelvis with US, CT, or MRI should be performed.
- Vulvar haematomas usually do not require surgical intervention, but in massive vulvar haematoma or haemodynamically unstable patients, surgical intervention, lavage and drainage is indicated.
- In vulvar laceration, suturing after conservative debridement is indicated with concomitant primary repair of any associated vaginal injuries.

Polytrauma, Damage Control and Mass Casualty Events

Urological trauma is often associated with significant and higher priority injuries in the polytraumatised patient. Damage

control principles govern the management of the severely injured patient and urologists need to understand their role in the context of polytrauma.

Damage control is a three-phase approach - rapid control of haemorrhage and wound contamination, resuscitation in the intensive care unit, and delayed definitive surgery.

Procedures should be directed at the rapid control of bleeding, debridement of dead and devitalised tissue, and minimising urinary extravasation by simple diversionary measures.

A mass casualty event is one in which the number of injured people is significantly higher than the number of healthcare providers available. Examples include the collapse of buildings or bridges, earthquakes, floods, tsunamis, train collisions, aircraft catastrophes, civilian terrorism.

Triage sorts patients into four groups:

1. Patients with life-threatening injuries that require immediate intervention, presenting with airway compromise, breathing failure and/or circulatory compromise from ongoing external haemorrhage.
2. Patients with severe but non-life-threatening injuries, in whom treatment can be acceptably delayed: major fractures, vascular injuries of the limbs and large soft tissue wounds.
3. 'Walking wounded' with minimal injuries.
4. Patients who are so severely injured that treatment would require allocation of resources and time that would deny other, more salvageable patients, timely care. These patients are given minimal or no treatment, and re-evaluated when resources become available. There is no absolute definition for this group because triage is

individualised according to the number and severity of casualties related to the available resources.

Principles for urological consultations to follow during a mass casualty scenario:

- Rule out under-triage by the surgeon in charge, and perform a rapid primary survey of every patient.
- Avoid unnecessary imaging procedures such as CT scans and retrograde urethrography. These procedures should be performed later, after mass casualty protocols have been suspended.
- Treat unstable patients who are to have surgery using damage control principles.
- Stable patients with suspected renal injuries should be transferred to the surgical ward without imaging procedures. Re-evaluate if there is any change in their haemodynamic status, or when possible as dictated by the constraints of the mass casualty event. Patients managed in this delayed fashion should be treated according to traditional trauma management protocols.
- 'Minimal acceptable' procedures should be performed in order to transfer patients to the surgical wards, e.g. suprapubic drainage of the bladder when bladder or urethral injuries are suspected, clamping and ligation of bleeding vessels from wounds to the external genitalia, etc.

This short booklet text is based on the more comprehensive EAU Guidelines (ISBN 978-94-92671-01-1) available to all members of the European Association of Urology at their website, <http://www.uroweb.org/guidelines>.